

CHARACTERIZATION OF THE LATE-VARISCAN I-TYPE GRANITIC ROCKS FROM THE WESTERN CARPATHIANS

I. BROSKA, I. PETRÍK, P. UHER

Geological Institute of the Slovak Academy of Sciences, Dúbravská cesta 9, 842 26 Bratislava, Slovak Republic

Two groups of the Variscan Carboniferous orogenic granitoids are found in the Western Carpathians area: the granitoids with affinity to the S-type and to the I-type. Both types were differentiated from tonalite up to granite (s.s.), granites prevailing in the S-type group and tonalites–granodiorites in the I-type. The S-type group possibly represents older collisional granitic suites (around 350 Ma by U–Pb and Rb–Sr dating), the I-type group is younger (around 305 Ma by U–Pb dating on zircons).

The I-type granitoids in the Western Carpathians are represented by the Sihla-type s.l. (Broska – Petřík 1993). The petrographical nature suggests relatively lower SiO₂ content (65–72 %) and increased contents of CaO, MgO and FeO. Also compatible trace elements have typically increased concentrations (Sr, Ba, Ni, V, Cr, Zr). Intense subsolidus alterations (chloritization and epidotization of biotite, sericitization of plagioclases) are characteristic of the I-type granitoids. The magnetite – allanite–sphene assemblage of accessory minerals associated with Mg – rich biotite indicate oxidation conditions in the melt. The zircon typology shows low alkalinity (index I.A 250 – 400) and intermediate to higher temperatures (index I.T > 350).

I-type granitoids were determined in the western part of Veporic unit, in the Modra massif (Malé Karpaty Mts.), Hlohovec body (Považský Inovec Mts.), Malá Fatra Mts., Trábeč Mts., Čierna Hora Mts.

Significant amount of oval shape mafic microgranular enclaves is locally present in the Western Carpathians I-type granitoids. We suppose that the enclaves are of magmatic origin and were quenched like globules of intermediate and basic composition in lower temperature felsic melt (in sense of Vernon 1983). This evokes an idea of contemporaneous basic and acid magmatism occurring during Variscan granitogenesis (Petřík – Broska 1989).

The genesis of Upper Carboniferous I-type granitoids might be connected with a renewed subduction regime which was suggested to occur on the southeastern flank of Gondwana (Finger – Steyrer 1990). Basic and intermediate magmas could have been products of subduction-related melting in the mantle wedge above subducted slab of the Paleotethys ocean floor. During emplacement of granite magmas operated extensional regime.

INDICATIONS OF CADOMIAN BACK ARC EXTENSION AND ACCRETION FROM THE NEOPROTEROZOIC OF EAST GERMANY

B. BUSCHMANN

Geological Institute, Freiberg Academy of Mining and Technology, B.-v.Cotta-Str. 2, D-09596 Freiberg, FRG

Within the Torgau–Doberlug–Synclinal Zone and extending into the southern rim of the Mid German Rise (the Southern Phyllitic Zone) a succession of Neoproterozoic to Middle Cambrian rocks occurs. Due to an extensive Cenozoic blanketing it is being investigated by numerous core-drills. The Neoproterozoic Rothstein Formation comprises: submarine basic volcanics with pillow intercalations & pyroclastic rocks / turbiditic greywacke–siltstone–mudstone sequences of a more distal turbiditic setting / pyritic black shales representing pelagic background sediments / hydrothermally altered varieties of the above mentioned rock types / hydrothermally exhalative cherts. This rock assemblage is indicative of an accumulation under distensive tectonic conditions typical for a back arc setting.

The metamorphic overprint of the Rothstein Formation increases from very low grade within the synclinoria to greenschist facies within the Southern Phyllitic Zone. Deformational features display striking similarities to structures developed within accretionary wedges (Knipe & Needham, 1986). These include downslope gravity driven deformations, deformations associated with accretion and collision–deformation related structures. Postcollision adjustments are difficult to record from drill cores as well as to tell from post–Cadomian tectonics.

Overstepping Lower and Middle Cambrian non-metamorphosed sediments accumulated under marine shallow water conditions. They are intruded by diabase sills and dykes of hitherto unknown age. Archeocyaths from Lower Cambrian limestones evidence Upper Atdabanian to Bottomian