

DEFORMATION OF GABBRO-AMPHIBOLITES IN THE SW BARRANDIAN ZONE (W BOHEMIAN MASSIF, NE BAVARIA) - ACADIAN TERRANE COLLISION

M. WEGER & L. MASCH

Institut für Mineralogie und Petrographie, Theresienstr. 41, 80333 Munich, Germany

The SW margin of the Barrandian zone, the Gabbro-Amphibolite Mass of Neukirchen-Kdyně (GAM), consists mainly of basic to intermediate intrusives of Cambrian age (511 ± 3 Ma, Gebauer & Grünenfelder, 1979; 504 ± 41 Ma, Miethig, A., 1993, in Köhler et al., 1993). To the West the Moldanubian (Mold)/Barrandian (Bar) border is marked by the E dipping Bohemian Pfahl and to the South by the NW dipping Central Bohemian Shear Zone. Both shear zones merge in the southwestern part of the GAM. Late Variscan granitoids intruded large parts of the Central Bohemian Shear Zone and Bohemian Pfahl of the Bohemian part of the GAM.




Fig. 1. Flinn diagram of 50 strain analyses in amphibolites of the Bavarian part of the GAM

Near the Mold/Bar border the basic and intermediate intrusives are metamorphosed and deformed to amphibolites. The Devonian age of the metamorphism is shown by amphibole K-Ar ages (380 Ma), whereas micas show later Variscan ages of 330 Ma. (Kreuzer et al., 1988). Devonian metamorphic ages are typical for the Acadian orogeny of the Appalachians. Since the Appalachian orogen of North America and the Variscan orogen of Europe belong to the same orogeny these ages should be called Acadian cooling ages. Bues (1992) determined medium pressure metamorphic conditions of 8–10 kb at 750° (lower granulite facies) at the border of the GAM decreasing to 6–7 kb at 680° (higher amphibolite facies) in the center of the Bavarian part of the GAM. This increase in grade of metamorphism towards the Mold/Bar border is consistent with the isograd pattern in the Bohemian part of the GAM. Towards the Northeast the metamorphic grade diminishes rapidly to virtually unmetamorphosed rocks of the Prague syncline.

The Acadian amphibole ages can be correlated with the lower granulite facies to higher amphibolite grade metamorphism. Towards the Mold/Bar border the basic intrusives, mostly gabbros, are successively deformed to mylonitic amphibolites. Undeformed metagabbros show typical magmatic microstructures and assemblages (pyroxene, olivine and large magmatic plagioclase, thermobarometry yielding submagmatic P,T). Low deformation leads to breakdown of olivine and pyroxene, growth of




Fig. 2a: 201 Mineral stretching lineations, Bavarian part of GAM, equal area net, lower hemisphere, 2–5–10–20%



Fig. 2b: 247 Poles to schistosity, Bavarian part of GAM, equal area net, lower hemisphere, 1–2–5–10–20%

large amphiboles and partial recrystallization of plagioclase. Mylonitic amphibolites consist of recrystallized and cataclastic amphiboles and recrystallized plagioclase domains showing core-mantle structures. Strains determined from the outlines of plagioclase domains show a rapid increase in deformation towards the Mold/Bar border. High strains are either prolate or oblate (Fig. 1) with mostly steep NE plunging mineral stretching lineation (Fig. 2a). The steep schistosity (60° – 90°) follows the Mold/Bar border from NW to E dip (Fig. 2b). Rare shear sense indicators are consistent with uplift of the GAM near the Mold/Bar border. The steeply NE plunging mineral lineation defines the direction of movement along the Mold/Bar border.

The degree of recrystallization of plagioclase can be correlated with strain values at sample and map scale and increases towards the Mold/Bar border. Steeply dipping schistosity, steeply NE plunging lineations, large pressure gradients, and shear sense indicators lead to the model of an Acadian collision between the Barrandian and Moldanubian terranes. Structures supporting the nappe model of Vollbrecht et al. (1989) based on DEKORP seismic profiles were not found.

The unusual geometry of the Bohemian Pfahl/Central Bohemian Shear Zone, one shear zone curving for 290° , can be explained by predeformational geometry of the edge of the Barrandian terrane. The collision lead to strong uplift and deformation of the edge of the SW Barrandian terrane. Acadian amphibole cooling ages in amphibolites with NE plunging mineral lineations are known from other parts of the western Bohemian massif, e.g. the Münchberg Gneiss Mass, the KTB deep drilling hole and the Zone of Erbdorf/Vohenstrauß. In the French Variscides Acadian movements were also recognized. In the Appalachians Acadian movements lead to the accretion of terranes along the continental margin (Rast, 1989; Keppie, 1993). Similar processes could have worked in the European Variscides, too. It is suggested, that Acadian movements and metamorphism are widespread in the European Variscides. They can only be traced in areas without major late Variscan plutonism, that probably destroyed many Acadian structures.

References:

- Bues, C. (1992): Dr. rer. nat. thesis, Munich Gebauer, D. & Grünfelder, M., 1979: Earth Planet. Sci. Lett. 42, 35–44
 Keppie, J.D. (1993): Geol. Rundsch. 83, 381–431
 Köhler, H. – Masch, L. – Miethig, A. – Pfeiffer, T. – Propach, G. – Weger, M. (1993): Eur. J. Min. 5 Beih. 2, 35–80
 Kreuzer, H. – Vejnar, Z. – Schüssler, U. – Okrusch, M., – Seidel, E. (1988): Proc. 1st. int. conf. Boh. Mass. Prague Rast, N., 1989: In: Bally, A.W. & Palmer, A.R., GSADNAG A, 323–348
 Vollbrecht, A. – Weber, K. – Schmoll, J. (1989): Tectonophys. 175,123–133

THE NORTH-WESTERN VARISCAN ODENWALD: HT-METAMORPHISM BENEATH AN ACTIVE CARBONIFEROUS BASIN

A.P. WILLNER¹ & A. KROHE²

¹ Institut für Mineralogie, Ruhr-Universität, D-44790 Bochum, Germany

² Institut für Mineralogie, Corrensstr.24, D-48149 Münster, Germany

The Variscan Odenwald is part of the Mid-German Crystalline Zone which partly represents mid-crustal levels of a magmatic arc. Around 360–330 Ma calcalkaline magmas synkinematically intruded along a major sinistral strike slip zone.

In the north-western Odenwald (Frankenstein area) abundant gabbros and diorites as well as few granodiorites intruded mainly amphibolites and minor metapelites. Prior to the intrusions these wall rocks were metamorphosed to regional amphibolite facies conditions at depth corresponding to only 1 to 2 kbar as it was shown by geothermobarometric studies of wall rocks using multivariant equilibria. Estimations of intrusion depth of the oldest gabbro intrusions correspond to this level. However, retrograde PT-data and estimation of intrusion depths of younger intrusions indicate an increase in pressure to maximal 3 kbar. Hence the PT-path was anticlockwise. Differential vertical movement was controlled by normal faulting concomitant with strike slip faulting. In the south-western prolongation of the strike-slip zone the Saar basin occurs with continuous sedimentation from Mid Devonian to Permian. From this configuration it is inferred that the major metamorphic imprint of the north-western Odenwald rocks occurred during subsidence underneath this basin. The high geothermal gradient results from extension as well as from voluminous accretion of basic magmas. A recent analogue is a pull-apart structure within a magmatic arc similar to the present Sunda arc.