


## Lamniporichnus vulgaris igen. et isp. nov.: traces of insect larvae in stone fruits of hackberry (*Celtis*) from the Miocene and Pleistocene of the Czech Republic



Lamniporichnus vulgaris igen. et isp. nov.:  
stopy hmyzích larev v peckách břestovce (*Celtis*)  
z miocénu a pleistocénu České republiky (Czech summary)

(1 text-fig., 1 plate)

RADEK MIKULÁŠ<sup>1</sup> – ZDENĚK DVOŘÁK<sup>2</sup> – †ILJA PEK<sup>3</sup>

<sup>1</sup> Geologický ústav Akademie věd České republiky, Rozvojová 135, 165 00 Praha 6, Czech Republic

<sup>2</sup> Severočeské doly, a.s., Doly Bílina, 418 29 Bílina, Czech Republic

<sup>3</sup> Přírodovědecká fakulta Univerzity Palackého, Svobody 26, 771 46 Olomouc, Czech Republic

Traces made probably by insect larvae in stone fruits of the Miocene hackberry, *Celtis lacunosa*, are described as *Lamniporichnus vulgaris* igen. et isp. nov. They were found in the opencast mine Merkur-North in the Most (North Bohemian) brown-coal basin. The newly proposed ichnogenus comprises borings in those wooden parts of fruits which contain a hollow filled with non-xylic digestible tissue (stones, nuts). The borings consist of a short cylindrical tunnel joining the stone (nut) surface and a natural hollow inside the xylic body. In comparison, the ichnogenus *Carporichnus* Genise, 1995 represents a longitudinal central cavity and a radial tunnel, both made by the tracemaker. Less favourably preserved material of *Celtis* sp. from the Pleistocene fill of karst cavities at Vitošov (Northern Moravia) provided also remains of presumably insect borings.

**Key words:** Insect borings, Miocene, Pleistocene, Most Basin, karst sediments, Czech Republic

### Introduction

The attention paid to insect traces in xylic substrates during the last decade has demonstrated that these traces are common in the fossil record since the Cretaceous (Genise 1995). However, as they had been usually overlooked before, systematic ichnology of them remains fragmentary.

The aim of the present contribution is to describe traces made probably by insect larvae in stone fruits of *Celtis lacunosa* (Reuss, 1861) Kirchner, 1957, from the Miocene of the Most (North Bohemian) brown-coal basin. Regardless the traces of comparable to identical morphology are quite common in Recent and presumably also in the Cenozoic fossil record, no ichnotaxonomical name has been given to them so far. The material from the North Bohemian Miocene is described herein as *Lamniporichnus vulgaris* igen. et isp. nov. Similar, but much less favourably preserved material of insect-damaged *Celtis* stone fruits is reported from the cemented fill of the karst cave at Vitošov (Northern Moravia).

### Geological settings

Merkur-North: Tertiary basins in Northwest Bohemia (Czech Republic) originated along the rift structure during the Eocene to Miocene. In the middle of the rift, the Most (North Bohemian) brown-coal basin is situated. In the Early Miocene, variable fluvial, limnic and swamp settings developed there. The past ecosystems are often well documented by the fossil flora, fish, insect, amphibian and reptile faunas (Kvaček – Fejfar 1993).

One of the rich fossil sites of the basin, which provided the fruit stones of *Celtis lacunosa* (Reuss) Kir-

chner bearing traces of insect larvae, is the opencast mine Merkur North of Chomutov. Here, the calcareous marls on the base of the so-called „Main Brown Coal Seam“ contain fossils of plants, terrestrial molluscs, insects and small mammals (Kvaček – Fejfar 1993). Stratigraphically, earlier part of the Early Miocene (Eggenburgian), lower Orleanium, mammal zone MN 3a is concerned. Reconstruction of the settings and vegetation (Kvaček – Fejfar 1993, Z. Kvaček *pers. comm.* 1998) show that shrubs to smaller trees of *Celtis lacunosa* were a part of forests that overgrew the brown coal swamps, together with *Acer tricuspidatum*, *Glyptostrobus europaeus*, *Alnus julianiformis*, with *Decodon* sp. in undergrowth, a.o.

The hackberry *Celtis lacunosa* was described first from freshwater limestones and travertines at Tuchořice (Reuss 1861). From the Most basin, few reports on its occurrence have been published so far. Holý in Čtyrský et al. (1964) first determined the species from the Merkur-North locality. Finds of endocarps of *C. lacunosa* from the layers overlaid by the Main Coal Seam are mentioned in the manuscript by Kvaček and Bůžek (1983). The insects at the Merkur-North locality are represented mostly by diving-beetles (J. Prokop, *pers. comm.* 1998).

Vitošov: The quarry at Vitošov exploits weakly metamorphosed limestones of the Devonian age. The quarrying led in 1997 to a discovery of karst cavities filled with breccia composed of biogenic particles and cemented with sinter. About 98 % of the breccia clasts are spherical bodies representing probably unfavourably preserved fruit stones of *Celtis* sp. Terrestrial molluscs are represented by *Granaria frumentum* (Draparnaud), *Clausilia* cf. *dubia* Drapanaud and fragments of several other taxa